

Volume 7  
Issue 1  
October 2022  
ISSN:5101195-3

# KOSOVA JOURNAL OF SURGERY



- Perspective: RIFAT LATIFI: On Being Minister of Health, Medical Diplomacy and the Transformation of Healthcare in Kosova
- SELMAN URANEUS: Damage Control Surgery in Severe Trauma
- JORG TEBAREK: Aortoiliac treatment with IVUS guidance as a standalone imaging solution: is it feasible and where are the benefits?
- JAMES HU: Familial Adenomatous Polyposis: Review of Current Diagnosis, Screening and Management



KOSOVA JOURNAL OF SURGERY  
Volume 7, Issue 1, October 2022  
ISSN:5101195-3  
info@koscs.org • www.koscs.org

**EDITOR-IN- CHIEF**

Rifat Latifi, MD, USA

**DEPUTY EDITORS-IN-CHIEF**

Luan Jaha, MD, Kosova  
Nexhmi Hyseni, MD, Kosova  
Ramadan Jashari, MD, Belgium  
Skender Topi, MD, Albania

**ASSISTANT EDITORS**

Fatjona Jahaj, Kosova

**EDITOR, BIostatISTICS**

Ilir Hoxha, MD, Kosova

**EDITOR OF SURGICAL HISTORY**

Isuf Dedushaj, MD, Kosova

**HONORARY EDITORS**

Sadri Bajraktari, MD, Kosova

**DESIGN AND LAYOUT**

Ardian Veliu, Kosova

This number is supported by  
Swiss Agency for Development and  
Cooperation SDC



© Copyright Kosovo Journal of Surgeons.  
All rights reserved. No parts of this magazine may be reproduced or transmitted in any form an by any means without written permission from Kosovo Journal of Surgeons. Kosovo Journal of Surgeons reserves the right to reject or accept any advertisement or advertisement material prior to publication.

**EDITORIAL BOARD**

**INTERNATIONAL**

Alban Neziri, MD, Switzerland  
Alberto R. Ferreres, MD,FACS Argentina  
Arben Baboci, MD, Albania  
Arben Beqiri, MD, Albania  
Arben Gjata, MD, Albania  
Artur Xhumani, MD, Albania  
Carlos Mesquita, MD, Portugal  
Demetrius Litwin, MD, FACS, FICS,USA  
Dritan Alushani, MD, Albania  
Emir Haxhija, MD, Austria  
Fatmir Dalladaku MD, Germany  
Georgious Tsoulfas, MD, FACS, FICS, Greece  
Gezim Dushaj, MD, Switzerland  
Girma Tefera, MD FACS, USA  
Giuseppe Nigri, MD, FACS, FRCS, Italy  
Hayato Kurihara, MD, FACS, Italy  
Horacio Rodriguez Rilo, MD, FACS, USA  
Itamar Askenazi, MD, Israel  
Joseph Bellal, MD, FACS, USA  
Jörg Tessarek, MD, Germany  
Kastriot Haxhirexha, MD,North Macedonia  
Kita Sallabanda, MD, Spain  
Lumnije Kqiku, DDs, Phd, Austria  
Maria Castaldi, MD, USA  
Minir Asani, MD, Germany  
Mustafa Ozsoy, MD, Turkey  
Moritz Felsenreich, MD, Austria  
Osman Dilek Nuri, MD, Turkey  
Peter Killcomons, MD, USA  
Philip Caushaj, MD, FACS, USA  
Ridvan Alimehmeti, MD, Albania  
Russel Andrews, MD, USA  
Roberto Bergamaschi, MD, FACS, USA  
Salomone Di Saverio MD, FACS, FRCS, England  
Selami Sozubir, MD, Turkey  
Selman Uranes, MD, FACS, Austria  
Tilsed Jonathan, MD, England  
Valon Baraliu, MD, Germany  
Xheladin Draçiqini, MD, Albania.

**NATIONAL**

Agreta Gashi- Gecaj, MD, Kosova  
Antigona Hasani, MD, Kosova  
Astrit Hamza, MD, Kosova  
Avdyl Krasniqi, MD, Kosova  
Aziz Mustafa, MD, Kosova  
Basri Lenjani, MD, Kosova  
Bedri Braha, MD, Kosova  
Belinda Pustina, MD, Kosova  
Besnik Bicaj, MD, Kosova  
Cen Bytyqi, MD, Kosova  
Dafina Mahmutaj, MD, Kosova  
Dukagjin Spanca, MD, Kosova  
Enver Fekaj, MD, Kosova  
Eqrem Shala, MD, Kosova  
Fadil Beka, MD, Kosova  
Fahredin Veselaj, MD, Kosova  
Faton Hoxha, MD, Kosova  
Fatos Sada, MD, Kosova  
Fehmi Ahmeti, MD, Kosova  
Ferat Sallahu, MD, Kosova  
Gani Çeku, MD, Kosova  
Harieta Zherka-Saraqini, MD, Kosova  
Hysni Jashari, MD, Kosova  
Isa Haxhiu, MD, Kosova  
Laura Leci, MD, Kosova  
Lulzim Vokri, MD, Kosova  
Nazmi Morina, MD, Kosova  
Nehat Baftiu, MD, Kosova  
Nexhat Rexha, MD, Kosova  
Osman Zhuri, MD, Kosova  
Raif Cavolli, MD, Kosova  
Sami Salihu, MD, Kosova  
Sejdi Statovci, MD, Kosova  
Shefqet Lulaj, MD, Kosova  
Shqiptar Demaci, MD, Kosova  
Sueda Latifi, MD, Kosova  
Talat Gjollli, MD, Kosova  
Vjollca Bince, MD, Kosova  
Xhevdet Cuni, MD, Kosova  
Xhevdet Tahiraj, MD, Kosova  
Ylber Zejnullahu, MD, Kosova

# KOSOVA COLLEGE OF SURGEONS

## KOSOVA JOURNAL OF SURGERY

October 2022 | Volume 7 | Issue 1 | www.koscs.org

## Contents



- 4** Perspective: **Rifat Latifi:** On Being Minister of Health, Medical Diplomacy and the Transformation of Healthcare in Kosova
- 13** **Selman Uraeus:** Damage Control Surgery in Severe Trauma
- 18** **Jorg TeBarek:** Aortoiliac treatment with IVUS guidance as a standalone imaging solution: is it feasible and where are the benefits?
- 25** **Lubomir Martinek:** Ileo-colic Anastomoses – A Review
- 32** **James Hu:** Familial Adenomatous Polyposis: Review of Current Diagnosis, Screening and Management
- 40** **Florin Iordache:** Updates in Fistula-in-Ano. A Narrative Review
- 56** **Selman Uranues, S.** Bile Duct Injury: Prevention Entails Analysis of Why it Happens, Not Only How it Happens?
- 63** **Abbas Smiley:** The Risk of Mortality in Geriatric Patients with Emergent Regional Enteritis is 12-fold Greater than that in Adult Patients: Female Sex, Hospital Length of Stay and Surgery As other Risk Factors of Mortality
- 82** **Adem Miftari:** Intravitreal Anti VEGF Application During Covid – 19 Pandemic in Kosova – Reviewer
- 88** **Abbas Smiley:** Longer stay in hospital is the risk factor for mortality whereas surgery is the protective factor in patients with blood in stool

# Ileo-colic Anastomoses – A Review

Assoc. Prof. Lubomir Martinek, MD., PhD. <sup>1</sup>

Katerina Vomackova, MD., PhD. <sup>2</sup>

<sup>1</sup>University Hospital and Faculty of Medicine Ostrava, Czech Republic

Consultant Surgeon, Head of Robotic Centre

President of Section of Endoscopic and Miniinvasive Surgery by Czech Surgical Society

<sup>2</sup>Olomouc University Hospital, Faculty of Medicine and Dentistry,

Palacky University Olomouc, Czech Republic

**Corresponding author:**

**LUBOMIR MARTINEK**

Odboje 1164, Vratimov, 739 32 Vratimov, Czech Republic

Phone: +420 603 500 627

E-mail: [lubomir.martinek@post.cz](mailto:lubomir.martinek@post.cz); [lubomir.martinek@osu.cz](mailto:lubomir.martinek@osu.cz); [lubomir.martinek@nnj.agel.cz](mailto:lubomir.martinek@nnj.agel.cz)

## Abstract

**Background:** Anastomosis between the small bowel and colon has been performed using a number of techniques. Despite a large number of trials and publications, a generally accepted optimal technique does not exist. There is a wide spectrum of procedures with variability in the technical details.

**Data sources:** The systematic literature search was performed by analysing the relevant databases – MEDLINE, EMBASE, Cochrane Library, and Google Scholar.

**Conclusions:** Guidelines regarding the ideal technical procedure for performing an ileocolic anastomosis are not currently available. Extracorporeal, stapled, side-to-side anastomosis with double-layer closure of the enterotomy is most commonly performed. Based on recent literature, laparoscopic/robotic right hemicolectomy with intracorporeal anastomosis currently seems to be the most promising technique.

**Keywords:** Ileo-colic anastomoses, techniques, review

## Introduction

An ideal ileocolic anastomosis has a minimal risk of leak, does not bleed, leads to a rapid recovery of bowel function, and stenosis does not occur in the long-term. It reduces the risk of recurrence in nonspecific inflammatory bowel diseases. In recent literature, the risk of leak in this location is reported to be 3-10%.<sup>1,2</sup> Bleeding and stenosis are relatively rare, and reliable data are lacking. The situation varies in cases of Crohn's disease, where stenosis usually develops as a consequence of disease recurrence with an incidence reaching up to 60%.<sup>3</sup>

Anastomotic complications are associated with a number of more or less established risk factors. For the surgeon, the most important factors are those he can influence, primarily the surgical technique.

Anastomosis between the small bowel and colon has been performed by various techniques. The open surgical procedure was expanded to a minimally invasive technique at the end of the 20<sup>th</sup> century. This currently includes a wide spectrum of procedures ranging from laparoscopically-assisted procedures and hand-assisted laparoscopic procedures to totally laparoscopic and

robotic operations. There is also variability in the technical details (hand-sewn anastomosis, combined stapled and hand-sewn anastomosis, completely stapled anastomosis), or in the configuration of the anastomosis (end-to-end, end-to-side, side-to-end, side-to-side, Kono-S anastomosis, isoperistaltic, or antiperistaltic anastomosis). Different types of sewing materials are used, continuous or interrupted sutures, and various types of staplers with varying staple height. Evaluation of blood perfusion using indocyanine green or hyperspectral imaging is rarely used.

Despite the number of published studies, a generally accepted optimal technique of performing an ileocolic anastomosis does not exist. Heterogeneity in patient sets in studies, as well as technical variability of the procedure, complicate objective comparison. This results in a lack of clearly formulated and generally accepted guidelines, based on robust data.

The aim of this review is not, and cannot be, a recommendation of one universal method. The work summarizes current comparisons of the most commonly used techniques of ileocolic anastomosis in terms of the surgical approach (open, laparoscopic, robotic) and basic technical details.

### **Open, Laparoscopic and Robotic Surgical Technique**

According to current data, laparoscopic right hemicolectomy offers a better postoperative course, lower risk of complications and similar oncological results as a conventional open procedure.<sup>4</sup> However, the quality of the laparoscopic complete mesocolic excision (CME) and the extent of lymph node dissection have been questioned.<sup>5,6</sup> The variability of blood supply to the right colon and the presence of critical structures (duodenum, venous trunk of Henle) makes the laparoscopic performance of CME technically demanding. Anania et al. reviewed 17 studies involving 2508 patients and reported that laparoscopic CME did not lead to a higher incidence of anastomotic leak when compared to the open approach, however, it was accompanied by a greater blood loss and higher frequency of conversion.<sup>7</sup> In 2021, the same group published a cohort of 5038 patients, confirming the benefits of a laparoscopic approach, especially in the early postoperative course with no difference in anastomotic complications.<sup>8</sup> Similar results are presented in other meta-analyses;<sup>9,10</sup> however, they often originate from specialized centres and their generalization is limited.

Robotic technique utilising a 3D view and instruments with EndoWrist® technology may reduce the technical difficulty of right hemicolectomy. According to the literature, the robotic approach is best at decreasing the number of conversions,<sup>11</sup> in the quality of CME,<sup>12</sup> and allows easier anastomosis construction<sup>11,13</sup> compared to laparoscopic surgery. Milone et al. did not find a difference in the frequency of leaks when comparing identical techniques of intracorporeal ileocolic anastomosis performed laparoscopically versus robotically.<sup>14</sup> Guadagni et al. in their analysis of 30 studies involving 2066 patients reported a significant reduction in anastomotic bleeding when using the robotic stapler (SmartClamp®Technology) compared to the laparoscopic stapler;<sup>15</sup> however, high quality RCT's with a sufficient number of patients are still lacking.

### **Extracorporeal vs. Intracorporeal Anastomosis**

There are two technical variants when performing the ileocolic anastomosis during laparoscopic right hemicolectomy— extracorporeal and intracorporeal. The intracorporeal technique has certain potential advantages. Bowel mobilisation in the abdominal cavity and traction to the mesentery are less extensive, the risk of unwanted bowel twisting is reduced, mesenteric defects are more easily closed, and it is possible to select the most advantageous area for removing the resected specimen. Extracorporeal anastomosis may be technically difficult in obese patients with a thick abdominal wall, or in patients with a broad or shortened mesentery. Disadvantages of an intracorporeal anastomosis are longer operating times, technical difficulty, and the potential risk of contamination when opening the bowel lumen inside the abdominal cavity.

Results of studies are not unanimous. Some authors have not confirmed significant differences between the two techniques.<sup>16,17</sup> Most recent works, including meta-analyses, report that patients with intracorporeal anastomosis have a more rapid functional recovery of the gastrointestinal tract, less pain, decreased morbidity, lower number of surgical site complications, shorter hospital stay and a better cosmetic results.<sup>18-22</sup> Cleary et al.<sup>23</sup> used propensity score matching to compare 379 intracorporeal and 650 extracorporeal anastomoses in a set of laparoscopically and robotically performed right hemicolectomies. In cases of intracorporeal anastomosis, longer operation time, shorter hospital stays, and less gastrointestinal and surgical site complications

were reported. In 2021, in a meta-analysis of 30 studies (1948 totally laparoscopically, 2369 laparoscopically-assisted), Zheng et al. presented a similar incidence of anastomotic complications, but a more rapid recovery of bowel function, less surgical site complications, and shorter hospital stay for patients with an intraabdominal anastomosis.<sup>24</sup> Hajinbandeh et al.<sup>22</sup> and Creavin et al.<sup>21</sup> used the same 4 RCT's in their meta-analysis (199 intracorporeal anastomoses, 200 extracorporeal anastomoses). They did not find a difference in the incidence of anastomotic complications, reported less postoperative pain and shorter hospitalization for the intracorporeal technique in the entire patient set, as well as in the subgroup of patients with cancer. There were fewer bleeding complications and a lower incidence of postoperative bowel obstruction at the cost of a longer operating time in patients with intracorporeal stapled side-to-side anastomosis. In addition, Creavin et al. also presented a significantly lower risk of postoperative gastrointestinal tract dysfunction in patients with intracorporeal anastomosis in their meta-analysis.<sup>21</sup>

### **Isoperistaltic vs. Antiperistaltic Ileocolic Anastomosis**

In the literature, not much attention is paid to the configuration of the ileocolic anastomosis. The antiperistaltic side-to-side anastomosis is referred to as a functional end-to-end anastomosis by some authors. Advantages include eliminating the risk of bowel twisting and a lower incidence of postoperative gastrointestinal tract dysfunction. However, it requires more extensive bowel mobilization than isoperistaltic anastomosis, which according to its supporters also limits the risk of twisting and eliminates possible traction. In isoperistaltic configuration, the enterotomy is closed by suture; the antiperistaltic configuration allows closure to be done by stapler. Recent randomized studies failed to establish that the configuration of the anastomosis influenced the incidence of leaks,<sup>25-27</sup> both have similar results in terms of safety, complications, and functions, including quality of life.<sup>26</sup>

### **Stapled and Hand-sewn Ileocolic Anastomosis**

In the literature, the stapler technique is often compared to hand-sewn ileocolic anastomosis. In a systematic review of 441 stapled and 684 hand-sewn ileocolic anastomoses presented in the Cochrane Database of Systematic Reviews from 2010 by authors Choy et al.,<sup>28</sup> it was reported that when the stapler was used, there was a

significantly lower incidence of leaks in the entire patient set (2.5% vs. 6.1%), as well as in the subgroup of patients operated for cancer (1.3% vs. 6.7%). The stapled anastomosis is recommended by authors as the standard for comparing anastomotic techniques. The following studies, however, led to differing results. Milone et al. did not find a difference between hand-sewn and stapled anastomosis in their work focussing primarily on enterotomy closure during laparoscopic or robotic intracorporeal side-to-side anastomoses.<sup>18</sup> The same risk of anastomotic leak (8.1%) in a set of 3208 patients was published in a prospective multicentric international audit by the European Society of Coloproctology collaborating group. When considering risk factors, the frequency of anastomotic leak in hand-sewn anastomoses was surprisingly significantly lower than when a stapler was used.<sup>29</sup> A higher risk of leaks for stapled ileocolic anastomoses (7.8%) was also reported by the Spanish national study ANACO (3.4%) including 52 centres comprising 1102 patients.<sup>30</sup> In addition, other recent multicentric studies state that the use of a stapler for a right hemicolectomy is a risk factor of anastomotic leak.<sup>29-32</sup> Nonetheless, generally, the stapled anastomosis is used demonstrably more often.<sup>29,30</sup>

The use of a linear stapler is associated with the need to close the enterotomy by optimal technique. Guadagni et al. evaluated the frequency of bleeding or leak with regard to the technique of enterotomy closure in robotic and laparoscopic right hemicolectomies. The authors did not confirm a significant difference between using a stapler or closing the enterotomy by a one-layer or a two-layer suture.<sup>15</sup> A different conclusion was reached by Milone et al. in their set of laparoscopically and robotically performed ileocolic anastomoses. Two-layer enterotomy closure led to a significantly lower risk of leak, with no difference in bleeding and with no stenosis.<sup>14</sup> The advantage of two-layer enterotomy closure was also confirmed by Reggio et al.<sup>33</sup>

Over-sewing the staple line is common in resections of the upper GI tract (esophagus, stomach), where the staple line is exposed to higher pressure. The usefulness of this step in the case of ileocolic anastomosis is ambiguous. Stapler manufacturers do not recommend over-sewing the staple line for risk of ischemia. In the already-mentioned prospective multicentric international audit of the European Society of Coloproctology collaborating group, over-sewing the staple line did not lead to better results<sup>32</sup> and similar conclusions have also been reached by other authors.<sup>34</sup>

## Ileocolic Anastomosis in Crohn's Disease

The issue of ideal ileocolic anastomosis in patients with Crohn's disease is slightly different. In addition to the above-mentioned complications, such as anastomotic leak, bleeding, or stenosis, there is also the risk of local disease recurrence. Although attention has long been paid to the technical performance of an ileocolic anastomosis in Crohn's disease, we still do not have an ideal technique and the presented results are controversial. A reduction in anastomotic leak in stapled side-to-side anastomoses compared to other types was confirmed by Similis et al.,<sup>35</sup> as well as according to a meta-analysis by authors Feng et al., with no difference in the frequency of recurrence.<sup>36</sup> Better results, including a reduction in the number of recurrences, were obtained after stapled side-to-side anastomosis compared to hand-sewn end-to-end anastomosis reported in the meta-analysis by He et al.<sup>37</sup> In a retrospective, multicentric observational study of the Italian Society of colorectal surgery focussing on patients with Crohn's disease (427 patients), Celentano et al. reported higher usage of the laparoscopic technique (72.8%), more frequent side-to-side configuration of the anastomosis (89%), and stapler preference (67%). Anastomotic leak was not dependent on the technique used for anastomosis.<sup>38</sup> Side-to-side configuration of the anastomosis helped reduce the risk of recurrence<sup>38</sup> and wide-lumen stapled anastomosis is recommended by certain surgical societies.<sup>39</sup> Contrarily, a better quality of life and lower frequency of re-hospitalizations are noted in end-to-end ileocolic anastomosis.<sup>40</sup> It has been established that neither staples nor non-absorbable suture material increase the risk of fistulas and recurrence.<sup>38</sup> Currently, the mesentery is studied as a potential source of recurrence. There are basically two types of surgical approaches: an extended mesentery resection proposed by authors Coffey et al.,<sup>3</sup> and the technique of side-to-side antimesenteric anastomosis described by Toru Kono in 2011 named the S-Kono anastomosis.<sup>41</sup> Robust data are still lacking.

## Discussion

Almost 30 years have passed since the first right hemicolectomy was performed by laparoscopic technique.<sup>42</sup> At first, only mobilization of the terminal ileum and colon were performed laparoscopically. Vessel ligation, resection, and anastomosis were performed openly via mini-laparotomy.<sup>43</sup> In 2004, Senagore et al.

described the technique of mediolateral mobilization, including vessel ligation, laparoscopically. The subsequent resection and extracorporeal anastomosis were performed by open procedure.<sup>44</sup> Technical difficulties with the extracorporeal anastomosis, especially in obese patients and those with a shortened mesentery, as well as efforts to minimize trauma to the abdominal wall, led to the development of a totally laparoscopic procedure. An intracorporeal laparoscopic ileocolic anastomosis was first mentioned by Roberto Bergamaschi in 1992<sup>45</sup> and has also been performed robotically since 2002.<sup>46</sup> Less frequent techniques of right hemicolectomy include hand-assisted laparoscopic resections; operations from one port or incision (SILS – Single Incision Laparoscopic Surgery); or procedures without mini-laparotomy, during which the resected specimen is usually extracted transvaginally (NOSE – Natural Orifice Specimen Extraction).

The minimally-invasive approach is generally characterized by a more favourable postoperative course with the same long-term and oncological results. The frequency of laparoscopically performed right hemicolectomies differs in the Western world. In Holland or Denmark, up to 75% of elective colorectal resection operations are performed laparoscopically. In other areas of Western Europe and in the USA, the frequency of laparoscopically performed hemicolectomies ranges between 30–50%.<sup>47,48</sup> An audit performed by the European Society of Coloproctology regarding right hemicolectomy stated that 54.6% of the procedures were performed laparoscopically.<sup>29</sup>

Isoperistaltic and antiperistaltic orientation of the anastomosis do not differ in safety or functionality (26). Although recent data question the superiority of a stapled anastomosis to hand-sewn, especially with regard to risk of leak,<sup>29-33</sup> a stapler is most frequently used during construction of the ileocolic anastomosis.<sup>29,30</sup> Intracorporeally performed ileocolic anastomosis also provides other potential benefits for the patient, such as less pain, more rapid recovery of bowel function, less complications, reduced surgical site infection rate, and better cosmetic effect.<sup>49-54</sup> This technique is not currently generally widespread. The situation is beginning to rapidly change with the use of the robotic approach. The robotic technique is safe, reduces the difficulty of a laparoscopic operation, reduces the number of conversions, and shortens the surgical training time. Short-term results of the robotic technique compared with laparoscopic right hemicolectomy are the same,<sup>55</sup> or in certain parameters

slightly better.<sup>56,57</sup> Long-term results are also comparable.<sup>58,59</sup> Disadvantages include high acquisition and operational costs of the robotic system. However, the number of studies is still low and robust data are lacking.

## Conclusions

Guidelines regarding the ideal technical procedure for performing an ileocolic anastomosis are not currently available, despite a number of published studies. Extracorporeal, stapled, side-to-side (isoperistaltic or antiperistaltic) anastomosis with double-layer closure of the enterotomy is most commonly performed.

Based on recent literature, laparoscopic/robotic right hemicolectomy with intracorporeal anastomosis currently seems to be the most promising technique. A number of interacting factors limit the reliability of results from randomized studies. Another method of data analysis (neural networks) could help find associations and support the surgeon in selecting the optimal technique for ileocolic anastomosis for each specific patient.

## Conflict of Interest Disclosure Statement

The author has no conflict of interest to declare.

## REFERENCES

1. Bakker IS, Grossmann I, Henneman D, et al. Risk factors for anastomotic leakage and leak-related mortality after colonic cancer surgery in a nationwide audit. *Br J Surg*. 2014; 101: 424-432.
2. Pellino G, Keller DS, Sampietro GM, et al. Italian Society of Colorectal Surgery SICCR (2020) Inflammatory bowel disease position statement of the Italian society of colorectal surgery (SICCR): Crohn's disease. *Tech Coloproctol*. 2020; <https://doi.org/10.1007/s10151-020-02183-z> [Epub ahead of print] Review)
3. Coffey CJ, Kiernan MG, Sahebally SM, et al. Inclusion of the Mesentery in Ileocolic Resection for Crohn's Disease is Associated With Reduced Surgical Recurrence. *J Crohns Colitis*. 2018 Nov 9;12(10):1139-1150.
4. Ding J, Liao GQ, Xia Y, et al. Laparoscopic versus open right hemicolectomy for colon cancer: a meta-analysis. *J Laparoendosc Adv Surg Tech. A* 2013;23(1):8-16.
5. Kileen S, Mannion M, Devaney A, et al. Complete mesocolic resection and extended lymphadenectomy for colon cancer: a systematic review. *Colorectal Dis*. 2014; 16(8): 577-594.
6. Pelz JOW, Wagner J, Lichthardt S, et al. Laparoscopic right-sided colon resection for colon cancer-has the control group so far been chosen correctly? *World J Surg Oncol*. 2018;16(1):117.
7. Anania G, Davies RJ, Bagolini F, et al. A. Right hemicolectomy with complete mesocolic excision is safe, leads to an increased

lymph node yield and to increased survival: results of a systematic review and meta-analysis. *Tech Coloproctol*. 2021;25(10):1099-1113.

8. Anania G, Arezzo A, Davies RJ, et al. A. A global systematic review and meta-analysis on laparoscopic vs open right hemicolectomy with complete mesocolic excision. *Int J Colorectal Dis*. 2021;36(8):1609-1620. 8.

9. Chaouch MA, Dougaz MW, Bouasker I, et al. Laparoscopic versus open complete mesocolon excision in right colon cancer: a systematic review and meta-analysis. *World J Surg*. 2019;43:3179-3190.

10. Alhassan N, Yang M, Wong-Chong N, et al. Comparison between conventional colectomy and complete mesocolic excision for colon cancer: a systematic review and pooled analysis: a review of CME versus conventional colectomies. *Surg Endosc*. 2019;33:8-18.

11. Yozgatli TK, Aytac E, Ozben V, et al. Robotic complete mesocolic excision versus conventional laparoscopic hemicolectomy for right-sided colon cancer. *J Laparoendosc Adv Surg Tech. A* 2019; 29(5):671-676.

12. Bae SU, Yang SY, Min BS. Totally robotic modified complete mesocolic excision and central vascular ligation for rightsided colon cancer: technical feasibility and mid-term oncologic outcomes. *Int J Color Dis*. 2019;34:471-479.

13. Spinoglio G, Bianchi PP, Marano A, et al. Robotic versus laparoscopic right colectomy with complete mesocolic excision for the treatment of colon cancer: perioperative outcomes and 5-year survival in a consecutive series of 202 patients. *Ann Surg Oncol*. 2018;25:3580-3586.

14. Milone M, Elmore U, Allaix ME, et al. Fashioning enterotomy closure after totally laparoscopic ileocolic anastomosis for right colon cancer: a multicenter experience. *Surg Endosc*. 2020; 34(2):557-563.

15. Guadagni S, Palmeri M, Bianchini M, et al. Ileo-colic intra-corporeal anastomosis during robotic right colectomy: a systematic literature review and meta-analysis of different techniques. *Int J Colorectal Dis*. 2021;36(6):1097-1110.

16. Feroci F, Lenzi E, Garzi A, et al. Intracorporeal versus extracorporeal anastomosis after laparoscopic right hemicolectomy for cancer: a systematic review and metaanalysis. *Int J Colorectal Dis*. 2013;28:1177-86.

17. Hellan M, Anderson C, Pigazzi A. Extracorporeal versus intracorporeal anastomosis for laparoscopic right hemicolectomy. *JSLs*. 2009;13(3):312-317.

18. Milone M, Elmore U, Vignali A, et al. Recovery after intracorporeal anastomosis in laparoscopic right hemicolectomy: a systematic review and meta-analysis. *Langenbecks Arch Surg*. 2018;403(1):1-10.

19. Ricci C, Casadei R, Alagna V, et al. A critical and comprehensive systematic review and meta-analysis of studies comparing intracorporeal and extracorporeal anastomosis in laparoscopic right hemicolectomy. *Langenbecks Arch Surg.* 2017;402(3):417-427.
20. Wu Q, Jin C, Hu T, et al. Intracorporeal versus extracorporeal anastomosis in laparoscopic right colectomy: a systematic review and meta-analysis. *J Laparoendosc Adv Surg Tech A.* 2017;27:348-357.
21. Creavin B, Balasubramanian I, Common M, et al. Intracorporeal vs extracorporeal anastomosis following neoplastic right hemicolectomy resection: a systematic review and meta-analysis of randomized control trials. *Int J Colorectal Dis.* 2021;36(4):645-656.
22. Hajibandeh S, Hajibandeh S, Mankotia R, et al. Meta-analysis of randomised controlled trials comparing intracorporeal versus extracorporeal anastomosis in laparoscopic right hemicolectomy: upgrading the level of evidence. *Updates Surg.* 2021;73(1):23-33.
23. Cleary RK, Kassir A, Johnson CS, et al. Intracorporeal versus extracorporeal anastomosis for minimally invasive right colectomy: A multi-center propensity score-matched comparison of outcomes. *PLoS One.* 2018;13(10):e0206277.
24. Zheng JC, Zhao S, Chen W, et al. Comparison of intracorporeal and extracorporeal anastomosis and resection in right colectomy: a systematic review and meta-analysis. *Langenbecks Arch Surg.* 2021;406(6):1789-1801.
25. Tarta C, Bishawi M, Bergamaschi R. Intracorporeal ileocolic anastomosis: a review. *Tech Coloproctol.* 2013;17:479-485.
26. Ibáñez N, Abrisqueta J, Luján J, et al. Isoperistaltic versus antiperistaltic ileocolic anastomosis. Does it really matter? Results from a randomised clinical trial (ISOVANTI). *Surg Endosc.* 2019;33(9):2850-2857.
27. Matsuda A, Miyashita M, Matsumoto S, et al. Isoperistaltic versus antiperistaltic stapled side-to-side anastomosis for colon cancer surgery: a randomized controlled trial. *J Surg Res.* 2015;196(1):107-112.
28. Choy PY, Bissett IP, Docherty JG, et al. Stapled versus handsewn methods for ileocolic anastomoses. *Cochrane Database Syst Rev.* 2011 Sep 7;(9):CD004320.
29. 2015 European Society of Coloproctology collaborating group. The relationship between method of anastomosis and anastomotic failure after right hemicolectomy and ileo-caecal resection: an international snapshot audit. *Colorectal Dis.* 2017. doi: 10.1111/codi.13646. Epub ahead of print.
30. Frasson M, Granero-Castro P, Ramos Rodriguez JL, et al. Risk factors for anastomotic leak and postoperative morbidity and mortality after elective right colectomy for cancer: results from a prospective, multicentric study of 1102 patients. *Int J Colorectal Dis.* 2016; 31: 105-114.
31. 2015 European Society of Coloproctology Collaborating Group Predictors for Anastomotic Leak, Postoperative Complications, and Mortality After Right Colectomy for Cancer: Results From an International Snapshot Audit. *Dis Colon Rectum.* 2020;63(5):606-618.
32. 2015 European Society of Coloproctology Collaborating Group. The impact of stapling technique and surgeon specialism on anastomotic failure after right-sided colorectal resection: an international multicentre, prospective audit. *Colorectal Dis.* 2018;20(11):1028-1040.
33. Reggio S, Sciuto A, Cuccurullo D, et al. Single-layer versus double-layer closure of the enterotomy in laparoscopic right hemicolectomy with intracorporeal anastomosis: a single-center study. *Tech Coloproctol.* 2015;19(12):745-750.
34. Fleetwood VA, Gross KN, Alex GC et al. Common side closure type, but not stapler brand or oversewing, influences side-to-side anastomotic leak rates. *Am J Surg.* 2017;213: 590-595.
35. Simillis C, Purkayastha S, Yamamoto T, et al. A meta-analysis comparing conventional end to-end anastomosis vs. other anastomotic configurations after resection in Crohn's disease. *Dis Colon Rectum.* 2007;50(10):1674-1687.
36. Feng JS, Li JY, Yang Z, et al. Stapled side-to-side anastomosis might be benefit in intestinal resection for Crohn's disease. *Medicine (United States).* 2018. <https://doi.org/10.1097/MD.00000000000010315>
37. He X, Chen Z, Huang J, et al. Stapled side-to-side anastomosis might be better than handsewn end-to-end anastomosis in ileocolic resection for Crohn's disease: a meta-analysis. *Dig Dis Sci.* 2014;59(7):15441551. h
38. Celentano V, Pellino G, Spinelli A, et al. Anastomosis configuration and technique following ileocaecal resection for Crohn's disease: a multicentre study. *Updates Surg.* 2021;73(1):149-156.
39. Bemelman WA, Warusavitarne J, Sampietro GM, et al. ECCO-ESCP consensus on surgery for Crohn's Disease. *J Crohns Colitis.* 2018;12(1):1-16.
40. Gajendran M, Bauer AJ, Buchholz BM, et al. Ileocecal anastomosis type significantly influences longterm functional status, quality of life, and healthcare utilization in postoperative Crohn's disease patients independent of inflammation recurrence. *Am J Gastroenterol.* 2018;113(4):576-583.
41. Kono T, Ashida T, Ebisawa Y, et al. A new antimesenteric functional end-to-end handsewn anastomosis: surgical prevention of anastomotic recurrence in Crohn's disease. *Dis Colon Rectum.* 2011;54(5):586-592.
42. Schlinkert RT. Laparoscopic-assisted right hemicolectomy. *Dis Colon Rectum.* 1991;34(11):1030-1031.
43. Young-Fadok TM, Nelson H. Laparoscopic right colectomy: five-step procedure. *Dis Colon Rectum.* 2000;43(2):267-271.

44. Senagore AJ, Delaney CP, Brady KM, et al. Standardized approach to laparoscopic right colectomy: outcomes in 70 consecutive cases. *J Am Coll Surg*. 2004;199(5):675-679.
45. Bergamaschi R, Haughn C, Reed JF 3rd, et al. Laparoscopic intracorporeal ileocolic resection for Crohn's disease: is it safe? *Dis Colon Rectum*. 2009;52(4):651-656.
46. Weber PA, Merola S, Wasielewski A, et al. Telerobotic-assisted laparoscopic right and sigmoid colectomies for benign disease. *Dis Colon Rectum*. 2002;45(12):1689-1694.
47. Alnasser M, Schneider EB, Gearhart SL, et al. National disparities in laparoscopic colorectal procedures for colon cancer. *Surg Endosc*. 2014;28(1):49-57.
48. Rea JD, Cone MM, Diggs BS, et al. Utilization of laparoscopic colectomy in the United States before and after the clinical outcomes of surgical therapy study group trial. *Ann Surg*. 2011;254(2):281-288.
49. Scotton G, Contardo T, Zerbinati A, et al. From laparoscopic right colectomy with extracorporeal anastomosis to robot-assisted intracorporeal anastomosis to totally robotic right colectomy for cancer: the evolution of robotic multi-quadrant abdominal surgery. *J Laparoendosc Adv Surg Tech A*. 2018;28(10):1216-1222.
50. Aiolfi A, Bona D, Guerrazzi G, et al. Intracorporeal versus extracorporeal anastomosis in laparoscopic right colectomy: an updated systematic review and cumulative meta-analysis. *J Laparoendosc Adv Surg Tech A*. 2020;30(4):402-412.
51. Saleh NB, Voron T, De'Angelis N, et al. Intracorporeal versus extracorporeal anastomosis in laparoscopic right hemicolectomy: results from the CLIMHET study group. *Tech Coloproctol*. 2020;24(6):585-592.
52. Bollo J, Turrado V, Rabal A, et al. Randomized clinical trial of intracorporeal versus extracorporeal anastomosis in laparoscopic right colectomy (IEA trial). *Br J Surg*. 2020;107(4):364-372.
53. Emile SH, Elfeki H, Shalaby M, et al. Intracorporeal versus extracorporeal anastomosis in minimally right colectomy: an updated systematic review and meta-analysis. *Tech Coloproctol*. 2019;23(11):1023-1035.
54. Martinek L, You K, Giuratrabocchetta S, et al. Does laparoscopic intracorporeal ileocolic anastomosis decrease surgical site infection rate? A propensity score-matched cohort study. *Int J Colorectal Dis*. 2018;33(3):291-298.
55. Solaini L, Cavaliere D, Pecchini F, et al. Robotic versus laparoscopic right colectomy with intracorporeal anastomosis: a multicenter comparative analysis on short-term outcomes. *Surg Endosc*. 2019;33(60):1898-1902.
56. Xu H, Li J, Sun Y, et al. Robotic versus laparoscopic right colectomy: a meta-analysis. *World J Surg Oncol*. 2014;12:27.
57. Ma Sa, Chen Y, Chen Y, et al. Short-term outcomes of robotic-assisted right colectomy compared with laparoscopic surgery: A systematic review and meta-analysis. *ASJSUR*. 2019;42(5):589-598.
58. Park JS, Kang H, Park SY, et al. Long-term oncologic after robotic versus laparoscopic right colectomy: a prospective randomized study. *Surg Endosc*. 2018;33(9):2975-81.
59. Kang J, Park YA, Baik SH, et al. A comparison of open, laparoscopic, and robotic surgery in the treatment of right-sided colon cancer. *Surg Laparosc Endosc Percutan Tech*. 2016;26(6):497-502.