




# KOSOVA JOURNAL OF SURGERY

## TRAUMA AND CRITICAL CARE SURGERY UPDATE: EXPANDING THE EVIDENCE — PART II



Milestones and Key Figures in the Development of Damage Control Surgery

VEDAT ELJEZI: State of the Art Review: Extracorporeal Membrane Oxygenation (ECMO): A Life-Saving Therapy And Its Challenges of Implementation In Emerging Healthcare Systems

NOAH ANTES, HUNTER ROW, THOMAS HALDIS, CORNELIUS DYKE: Transcatheter Aortic Valve Replacement for Severe Aortic Stenosis: A Review and Special Considerations

LULEJETA ATLANTIDA LATIFI, RIFAT LATIFI: Damage Control Surgery: An Update

WADE HOPPER, MENTOR AHMETI: State of the Art Interventions in Geriatric Trauma

DANIEL SCHEESE, TRAVIS SULLIVAN, JEFFREY HAYNES, EDGAR RODAS: Navigating the Challenges of Managing Pediatric Trauma: Current Trends, Evidence, and Practical Pathways

BELINDA DE SIMONE, FAUSTO CATENA, RIFAT LATIFI: Venous Thromboembolism Prophylaxis in Trauma and Critically Ill Patients: A State-of-the-Art Evidence-Based Review

OLUWAFEMI P. OWODUNNI, ELLIOTT HAUT: Implementing Sustainable Systems for Achieving Excellence in Venous Thromboembolism Prevention

**Volume 10, Issue 1, Part II:  
Trauma and Critical  
Care Surgery Update:  
Expanding the Evidence  
April 2026  
ISSN: 3027-5008 (Online)  
ISSN: 3027-5016 (Print)**



# Implementing Sustainable Systems for Achieving Excellence in Venous Thromboembolism Prevention

Oluwafemi P. Owodunni, MD, MPH

Elliott R. Haut, MD, PhD, FACS

Department of Emergency Medicine, the University of New Mexico Hospital,  
Albuquerque, NM (OPO)

Division of Acute Care Surgery, Department of Surgery (ERH)  
Department of Anesthesiology and Critical Care Medicine (ERH)

Department of Emergency Medicine (ERH)

The Johns Hopkins University School of Medicine, Baltimore, Maryland

The Armstrong Institute for Patient Safety and Quality (ERH)

Johns Hopkins Medicine, Baltimore, Maryland

Department of Health Policy and Management (ERH)

The Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland

oowodunni@salud.unm.edu; ehaut1@jhmi.edu

## Corresponding Author:

## ELLIOTT R. HAUT, MD, PHD, FACS

Vice Chair of Quality, Safety & Service, Department of Surgery

Professor of Surgery, Anesthesiology / Critical Care Medicine (ACCM) and Emergency Medicine

Division of Acute Care Surgery, Department of Surgery

The Johns Hopkins University School of Medicine

Professor of Health Policy & Management

The Johns Hopkins University Bloomberg School of Public Health

Core Faculty, The Armstrong Institute for Patient Safety and Quality, JHM

Sheikh Zayed 6107C; 1800 Orleans St.; Baltimore, MD 21287

*Email:* ehaut1@jhmi.edu (email)

@elliottthaut (Twitter)

Haut (<https://orcid.org/0000-0001-7075-771X>)

## Abstract

Venous thromboembolism (VTE) is a common preventable cause of morbidity and mortality in hospitalized

patients. Despite robust guidelines and decades of data, there is still underperformance in three important aspects of defect-free VTE prevention. The objectives



of this perspective are threefold: first, to delineate the critical steps of risk-appropriate VTE prevention i.e., 1) risk assessment (balancing risks of VTE and bleeding), 2) prescription of risk-appropriate prophylaxis, and 3) reliable administration of all prescribed VTE prophylaxis doses. Second, to examine why existing systems fail to deliver consistent risk-appropriate care; and third, to propose a pragmatic and scalable accountability framework to close these gaps. Drawing from evidence and implementation science, we propose a system-wide strategy that integrates electronic clinical decision support, individualized clinician feedback, and targeted nurse-specific and patient-centered education to reduce missed doses. Sustained success will require institutions to integrate the evidence-based processes within system-level infrastructure, with ongoing surveillance and quality improvement.

**Key words:** Deep Vein Thrombosis, Pulmonary Embolism, Venous Thromboembolism, Quality Improvement, Patient Safety, Thromboprophylaxis

## Introduction

Venous thromboembolism (VTE) comprising deep venous thrombosis (DVT) and pulmonary embolism (PE), is a serious and often fatal complication experienced by hospitalized patients.<sup>1,2</sup> Surgical patients, especially those with trauma, orthopedic surgery, or cancer, are among the highest-risk populations. The American Heart Association estimates that recent surgery accounts for approximately one-quarter of VTE events, with highly heterogeneous risk across demographic and clinical factors.<sup>2,3</sup>

Across the surgical population, symptomatic VTE is estimated at 1.6% in the early postoperative period, and as many as 56% of events are first detected following discharge.<sup>2</sup> Rates are estimated to be higher for emergency and some nonemergent procedures alike, and may be attributable to approach (laparoscopic versus open), duration and complexity.<sup>2,4,5</sup> The procedure-level data illustrate the gradient in VTE variation based on surgery type, for example, <0.1% after laparoscopic cholecystectomy, ≤4% after open small-bowel resection and ≤10% following emergency open total proctocolectomy.<sup>4,5</sup> The VTE risk after oncologic procedures is amplified by advanced staging and nonminimally invasive approach.<sup>6</sup> Additional risk factors include age, body mass index, previous VTE, reduced mobility, cardiopulmonary diseases,<sup>7-9</sup> and other clinical parameters.<sup>9</sup>

Prior to the implementation of routine VTE prophylaxis the downstream morbidity was substantial.<sup>2,10</sup> Post-thrombotic syndrome was noted in 30–50% of patients after proximal DVT, and chronic thromboembolic pulmonary hypertension in <4% of PE survivors within the first two years.<sup>2,10</sup> The burden on healthcare systems also aligns with the disability burden. Higher readmission rates and extended hospital stay have been associated with VTE, leading to increased resource utilization.<sup>11,12</sup> In the context of financial implications, the economic footprint is large. The direct medical costs are approximated at \$12,000–\$15,000 per patient in the first year and ≥\$23,000 following complications.<sup>2,10</sup> The overall estimated cost for VTE in the United States is \$7–10 billion.<sup>2</sup>

## Current Guideline Consensus and Areas of Controversy

### Consensus

Major societies including the American Society of Hematology, American College of Chest Physicians, and National Institute for Health and Care Excellence provide guidance on several key principles for VTE prevention in surgical patients.<sup>1,3,4,8</sup> This guidance converges on the need to stratify, balance measures between prophylaxis and bleeding risk and consider extended prophylaxis when post-discharge risk remains elevated.<sup>1,3,4,8</sup>

Individualized risk assessment using validated tools such as Caprini, Rogers, or a “bucket model” can help translate procedure class and patient factors into an actionable risk stratum.<sup>1,3,4,8</sup> For patients in the moderate or high-risk strata without a prohibitive bleeding profile, pharmacologic prophylaxis is routinely recommended.<sup>1,3,13</sup> When anticoagulation is contraindicated or deferred for immediate postoperative concern, mechanical prophylaxis using sequential compression devices is recommended. Importantly, the effectiveness of these devices is tied to adherence in use.<sup>1,3</sup> Combination therapy with pharmacologic and mechanical prophylaxis is appropriate for patients at the highest risk stratum when bleeding risk is acceptable.<sup>1</sup> Additionally, extended prophylaxis typically for up to 4 weeks is also recommended in surgical patients<sup>1,3,4</sup> similar to nonsurgical patients.<sup>14</sup>

### Areas of Controversy

Despite broad consensus, meaningful gaps remain. There is a commonly held misbelief that ambulation alone provides effective VTE prophylaxis. Early mobilization remains desirable for recovery and cardiopulmonary



benefits, but it is not an adequate stand-in for pharmacologic or mechanical prophylaxis. Patients should not have pharmacologic VTE prevention withheld or discontinued solely because they are ambulating, absent of a competing contraindication.<sup>15</sup> Subspecialty risk stratification also remains debated with limited head-to-head comparisons.<sup>3,4,8,16,17</sup> Except for orthopedic and oncology, the ideal duration of prophylaxis is poorly defined and real-world adherence to extended prophylaxis varies.<sup>18–20</sup> Furthermore, unfractionated heparin or low-molecular weight heparin have been the mainstay of prophylaxis in hospitalized patients. However, newer data on the use of aspirin for prophylaxis is spreading from patients undergoing orthopedic joint replacement to other populations such as trauma.<sup>21,22</sup> The data from the PREVENT CLOT pragmatic, multicenter, randomized noninferiority trial, conducted in 21 trauma centers across North America demonstrated that aspirin was noninferior to low-molecular weight heparin for preventing mortality and symptomatic VTE in patients with a fracture. These findings support risk-appropriate, patient-specific strategies rather than universal heparin-based regimens.<sup>21</sup> Aspirin has been used for decades for VTE prophylaxis after joint replacement surgery. This paper now shows that aspirin is also effective for VTE prevention in the orthopedic trauma population.

### **System-Level Interventions Highlighting the Evidence and Impact**

#### *Risk-appropriate Prescribing Practices*

The Johns Hopkins collaborative improved VTE prophylaxis by incorporating risk assessment and tools to assist frontline clinicians with prescribing VTE prophylaxis into everyday workflow.<sup>23</sup> Starting in 2005, a multidisciplinary team designed a paper checklist and then transitioned to a mandatory computerized provider order entry (CPOE) decision support tool that encouraged completion of risk stratification embedded within admission orders.<sup>23,24</sup> Early initiatives exposed barriers to adoption, informing transition to a decision support tool with auto-populated patient data, default risk-appropriate prophylaxis guidance, and a true mandatory function.<sup>23</sup> Performance was monitored via a database with routine feedback to services and individual clinicians. This full compendium was coupled with targeted education and local champions to maintain momentum. This effort resulted in a stepwise improvement from paper to electronic implementation i.e., risk-appropriate prophylaxis

prescribing increased to 80% in surgery subspecialties and 92% on medicine services by 2011, and continuous improvements after dedicated feedback cycles were established.<sup>23</sup>

In a level-1 trauma service of the same hospital, the same CPOE-embedded decision support tool was evaluated. Risk-stratification documentation within 24 hours increased from 3.0% at baseline to 97.8% after implementation.<sup>25</sup> Risk-appropriate prophylaxis prescribing increased from 66.2% to 84.4%, with the largest improvements observed among patients without contraindications. Furthermore, preventable VTE significantly decreased from 1.0% to 0.17%.<sup>25</sup> Individualized performance feedback was provided to general surgery residents in the context of their individual VTE-prophylaxis prescribing practices. The results were robust, showing significant improvement in appropriate prophylaxis for patients admitted by general surgery residents, increasing from 78.1% to 93.9%.<sup>26</sup> A beneficial unintended consequence of this intervention was observed among advanced practice providers (nurse practitioners and physician assistants). Despite receiving no feedback, they significantly improved in their risk-appropriate prescribing practices from 75.1% to 84.9%.<sup>26</sup>

#### *Frontline Nurses' Perspectives, Administration Practices and Education*

The CPOE had unquestionably improved prescription of venous thromboembolism prophylaxis, yet a completed order did not guarantee a risk-appropriate dose was administered to the patient.<sup>27</sup> Patients were not receiving their prescribed doses of prophylaxis medications, even when prescribed appropriately. At the bedside, execution faltered for pragmatic and cultural reasons i.e., patients decline injections, ambulation is incorrectly considered as a waiver, transient clinical changes prompting ad hoc holds and competing tasks or handoffs that disrupted workflow. An early refusal may often cascade into inertia, with subsequent doses occasionally not offered. Additionally, documentation euphemisms such as “off unit” or “patient refused” can obscure true missed opportunities and unit-level cultures drove much of the variation in administration.<sup>27</sup>

We tackled this problem head-on by working with patients and other key stakeholders to create interventions to improve missed doses of prophylaxis in hospitalized patients. A cluster-randomized trial tested whether nurse-focused education, created and imple-

mented with buy-in from nurse leadership, could close the gap between risk-appropriate prophylaxis prescribing and administration practices.<sup>28</sup> There were improvements after implementation of the nurse-focused education module, with significant increases in administration of prescribed risk-appropriate VTE prophylaxis.<sup>28</sup> Our online nurse education module is freely available online via a collaboration with the Anticoagulation Forum (<https://acforum.org/VTE-C1/story.html>).

### *Missed Doses Remain a Pervasive Issue*

Across hospitals a persistent gap remains between ordering and delivery, turning a preventable event into a predictable one.<sup>29</sup> At our institution, an audit revealed that 12% of prescribed doses of pharmacologic prophylaxis were not given and nearly 60% of these doses not administered were attributable to patient or family refusal.<sup>29</sup> Such refusals are often multifactorial i.e., may reflect transient discomfort, concern about injections or bruising, misconceptions about ambulation and uneven bedside communication.<sup>27-30</sup> Regardless of the cause, the clinical end point is identical as high-risk patients will remain at risk of VTE. Reframing nonadministration as a modifiable delivery failure rather than a fixed patient preference moved the attention from individual blame to system opportunities. We developed a patient-centered education bundle, co-developed by individual patients and organizational support from the National Blood Clot Alliance, providing opportunities for targeted patient communication and reliable bedside workflows that can convert more prescribed risk-appropriate prophylaxis doses into administered doses.

In a controlled implementation effort, we leveraged information technology within the electronic medication administration record that notified a dedicated health educator whenever a risk-appropriate prescribed prophylaxis dose was not given.<sup>30</sup> The educator first clarified the reason with the bedside nurse; if refusal was the driver, the patient received a brief one-time education bundle tailored to their preference including 1) a face-to-face conversation, 2) a two-page multilingual handout, and/or 3) a 10-minute video on a tablet. Content and delivery strategies were created and informed by a modified Delphi process with input from national thrombosis organizations, patient stakeholders, and a patient and family advisory council.<sup>30,31</sup> Despite the educator's part-time presence and delivery to only about one-third of eligible encounters, risk-appropriate VTE prophylaxis administration

significantly improved. On intervention units, overall nonadministration significantly reduced from 9.1% to 5.6%, patient refusal decreased from 5.9% to 3.4%, and missed doses for other reasons also decreased from 2.3% to 1.7%.<sup>30</sup>

### *Mitigating Bias: An Unintended Consequence of Quality Improvement Initiatives*

An important lesson from our implementation efforts is that robust, system-level quality improvement can mitigate and eliminate preexisting disparities while improving the standard of care at the bedside.<sup>32-34</sup> The mandatory CPOE increased overall adherence on surgical and medical services, and it also mitigated race-based disparities in risk-appropriate VTE prescribing.<sup>32</sup> Although it was not our primary goal, making the correct choice the default yielded a consequential byproduct of standardized care. Furthermore, subgroup analyses showed that the real-time patient-centered education bundle had comparable reductions in missed risk-appropriate VTE prophylaxis doses across race and sex groups, countering concerns that a technology-enabled, educator-mediated approach may preferentially be advantageous to certain groups.<sup>33,34</sup> Together, these studies highlight two important points. First, disparities are multifactorial and arise from defects across the care continuum i.e., variable clinician judgment at ordering and inconsistent bedside communication during administration. Second, a multidisciplinary, stakeholder-engaged design can mitigate disparities while improving care for all patients.

### *Success, Scaling, Dissemination, and Implementation Efforts*

The successful implementation of these interventions at one academic center in our health system was achieved when three conditions were met: 1) visible leadership accountability (clear buy-in at the service and executive levels), 2) health informatics capacity, and 3) unit-based engagement (charge-nurse stewardship and bedside champions).<sup>35,36</sup> These interventions have subsequently been disseminated to a community hospital in our health system with comparable reductions in missed and refused doses.<sup>36,37</sup> These interventions were implemented at nine trauma centers that are part of the Coalition of Leaders in Thromboembolism study group (CLOTT-3), supported by the Patient-Centered Outcomes Research Institute and run through the Coalition for National Trauma Research.

## Opportunities in Low- and Middle-Income Countries (LMICs)

Global inequities in VTE prevention are multifactorial and represent differences in healthcare infrastructure, economic resources, and slow adoption of evidence-based guidelines.<sup>38</sup> In a Vietnamese study, Nguyen and colleagues found that clinicians who understood the risk but underestimated the benefit, and failure to operationalize institutional protocols, were key barriers to the optimal adoption of VTE prevention strategies.<sup>39</sup> In contrast, a European cancer center demonstrated that risk-stratified protocols increased compliance to 91%, reducing postoperative VTE by nearly 80% and sustaining that success for years.<sup>40</sup> While these studies were performed in different settings and with varying institutional capacity, defect-free VTE prevention can be replicated globally. While local resources may differ in different countries, the pillars i.e., risk assessment, education, and operationalizing care do not. Moreover, training, stewardship, and protocols should be the minimum standard of patient safety individualized to match institutional capacity. VTE prophylaxis is cost effective and many of the medications routinely used are relatively inexpensive (i.e. heparin, aspirin).

In summary, our shared architecture is straightforward and can be adopted in different settings. First, defect-free care was a single shared aim which set the expectation and anchored measurement.<sup>41</sup> Second, real-time data capacity must be delivered to the people who can act on it (i.e., nurses see real-time nonadministration alerts, prescribers receive succinct performance reports, and unit leaders track compliance).<sup>35</sup> Third, persistent myth-busting is built into scripts and huddles, especially the misconception that ambulation obviates anticoagulation and the habit of converting a one-time refusal into a standing hold.<sup>15,35</sup> Sustainability followed from making the right action the path of least resistance. We used an implementation toolkit including standardized nurse education, and patient-education materials (freely available online in 13 languages at <https://www.hopkinsmedicine.org/armstrong-institute/improvement-projects/vte>), posters, badge cards for nurses, and monthly feedback loops embedded in routine leadership rounds.

## Conclusion

We present our experience implementing patient-centered interventions directed at the most defect-prone areas in VTE prevention. Across all settings, whether at a tertiary

or a community setting in high resourced or LMICs, the principles do not change. The evidence is strong enough to demand action in the context of performing standardized risk assessments, prescribing risk-appropriate prophylaxis, and ensuring every ordered dose is administered. When prevention is integrated into workflow, reinforced by culture, and upheld through accountability, the outcome transcends abstract improvement and becomes a tangible reduction of preventable VTE and the reaffirmation of what reliable care should look like.

**Author Contributions:** Dr. Owodunni and Dr. Haut contributed equally to all phases of article development including concept, drafting, critical revisions, and approval of the final version of the manuscript. Both authors meet authorship criteria per ICMJE recommendations.

**Funding:** No funding was obtained for this article.

### Conflicts of Interest:

Dr. Owodunni reports no conflicts of interest or financial disclosures.

Dr. Haut reports research funding from The Patient-Centered Outcomes Research Institute (PCORI) and the Agency for Healthcare Research and Quality (AHRQ). Dr. Haut is an unpaid member of the Board of Directors of the National Blood Clot Alliance (NBCA) and the Coalition for National Trauma Research. Dr. Haut is the current Editor-in-Chief of *Trauma Surgery & Acute Care Open*.

**Data Availability:** Not applicable. No original data reported.

**Consent:** No new or original patient data are reported in this article and thus patient consent is not necessary or applicable.

**Ethical Approval:** Ethical approval is not needed for this paper since no new or original patient data are reported in this article.

**Pre-Registration:** Not applicable to this article.

## References:

1. Anderson DR, Morgano GP, Bennett C, et al. American Society of Hematology 2019 guidelines for management of venous thromboembolism: prevention of venous thromboembolism in surgical hospitalized patients. *Blood Adv.* 2019;3(23):3898-3944. doi:10.1182/bloodadvances.2019000975
2. Henke PK, Kahn SR, Pannucci CJ, et al. Call to Action to Prevent Venous Thromboembolism in Hospitalized Patients: A Policy Statement From the American Heart Association. *Circulation.* 2020;141(24):e914-e931. doi:10.1161/CIR.0000000000000769

3. Gould MK, Garcia DA, Wren SM, et al. Prevention of VTE in Nonorthopedic Surgical Patients. *Chest*. 2012;141(2 Suppl):e227S-e277S. doi:10.1378/chest.11-2297
4. Patel SV, Liberman SA, Burgess PL, et al. The American Society of Colon and Rectal Surgeons Clinical Practice Guidelines for the Reduction of Venous Thromboembolic Disease in Colorectal Surgery. *Diseases of the Colon & Rectum*. 2023;66(9):1162. doi:10.1097/DCR.0000000000002975
5. Lavikainen LI, Guyatt GH, Sallinen VJ, et al. Systematic Reviews and Meta-analyses of the Procedure-specific Risks of Thrombosis and Bleeding in General Abdominal, Colorectal, Upper Gastrointestinal, and Hepatopancreatobiliary Surgery. *Ann Surg*. 2024;279(2):213-225. doi:10.1097/SLA.0000000000006059
6. Theochari CA, Theochari NA, Mylonas KS, et al. Venous Thromboembolism Following Major Abdominal Surgery for Cancer: A Guide for the Surgical Intern. *Curr Pharm Des*. 2022;28(10):787-797. doi:10.2174/1381612828666220217140639
7. Xia ZH, Chen WH, Wang Q. Risk factors for venous thromboembolism following surgical treatment of fractures: A systematic review and meta-analysis. *Int Wound J*. 2023;20(4):995-1007. doi:10.1111/iwj.13949
8. Tikkinen KAO, Siegal DM, Devereaux PJ, et al. The CLUE postsurgery VTE risk instrument for abdominal and pelvic surgery: validation of patient risk factor component. *Blood Adv*. 2025;9(15):3837-3844. doi:10.1182/bloodadvances.2024015515
9. Pastori D, Cormaci VM, Marucci S, et al. A Comprehensive Review of Risk Factors for Venous Thromboembolism: From Epidemiology to Pathophysiology. *Int J Mol Sci*. 2023;24(4):3169. doi:10.3390/ijms24043169
10. Tipirneni KE, Bauter L, Arnold MA, Audlin JA, Ryan J, Marzouk M. Association of Prolonged-Duration Chemoprophylaxis With Venous Thromboembolism in High-risk Patients With Head and Neck Cancer. *JAMA Otolaryngol Head Neck Surg*. 2021;147(4):320-328. doi:10.1001/jamaoto.2020.5151
11. Mallick S, Aiken T, Varley P, et al. Readmissions From Venous Thromboembolism After Complex Cancer Surgery. *JAMA Surg*. 2022;157(4):312-320. doi:10.1001/jamasurg.2021.7126
12. Dobaria Bs V, Hadaya J, Ebrihiminan S, Verma A, Sanaiha Y, Benharash P. Impact of Venous Thromboembolism on Readmissions and Resource Use Following Emergency General Surgery. *Am Surg*. 2022;88(10):2436-2439. doi:10.1177/00031348221101478
13. Bartlett MA, Mauck KF, Stephenson CR, Ganesh R, Daniels PR. Perioperative Venous Thromboembolism Prophylaxis. *Mayo Clin Proc*. 2020;95(12):2775-2798. doi:10.1016/j.mayocp.2020.06.015
14. Spyropoulos AC, Goldin M, Ageno W, et al. Rivaroxaban Plus Aspirin for Extended Thromboprophylaxis in Acutely Ill Medical Patients: Insights from the MARINER Trial. *TH Open*. 2022;6(3):e177-e183. doi:10.1055/s-0042-1750379
15. Lau BD, Murphy P, Nastasi AJ, et al. Effectiveness of ambulation to prevent venous thromboembolism in patients admitted to hospital: a systematic review. *CMAJ Open*. 2020;8(4):E832-E843. doi:10.9778/cmajo.20200003
16. Khatri A, Machin M, Vijay A, Salim S, Shalhoub J, Davies AH. A Review of Current and Future Antithrombotic Strategies in Surgical Patients-Leaving the Graduated Compression Stockings Behind? *J Clin Med*. 2021;10(19):4294. doi:10.3390/jcm10194294
17. Kakkar AK, Cohen AT, Tapson VF, et al. Venous thromboembolism risk and prophylaxis in the acute care hospital setting (ENDORSE survey): findings in surgical patients. *Ann Surg*. 2010;251(2):330-338. doi:10.1097/SLA.0b013e3181c0e58f
18. Aiken TJ, King R, Russell MM, Regenbogen SE, Lawson E, Zafar SN. Venous thromboembolism prophylaxis following colorectal surgery: a survey of American Society of Colon and Rectal Surgery (ASCRS) member surgeons. *J Thromb Thrombolysis*. 2023;55(2):376-381. doi:10.1007/s11239-022-02733-2
19. Logan CD, Hudnall MT, Schlick CJR, et al. Venous Thromboembolism Chemoprophylaxis Adherence Rates After Major Cancer Surgery. *JAMA Netw Open*. 2023;6(9):e2335311. doi:10.1001/jamanetworkopen.2023.35311
20. Nemeth B, Lijfering WM, Nelissen RGHH, et al. Risk and Risk Factors Associated With Recurrent Venous Thromboembolism Following Surgery in Patients With History of Venous Thromboembolism. *JAMA Netw Open*. 2019;2(5):e193690. doi:10.1001/jamanetworkopen.2019.3690
21. Major Extremity Trauma Research Consortium (METRC). Aspirin or Low-Molecular-Weight Heparin for Thromboprophylaxis after a Fracture. *New England Journal of Medicine*. 2023;388(3):203-213. doi:10.1056/NEJMoa2205973
22. Leyva H, Fanikos J, Lan Z, et al. Aspirin for Venous Thromboembolism Prophylaxis in Hip Fracture, Total Knee, and Hip Arthroplasty. *The American Journal of Medicine*. 2025;138(9):1185-1196. doi:10.1016/j.amjmed.2025.03.020
23. Streiff MB, Carolan HT, Hobson DB, et al. Lessons from the Johns Hopkins Multi-Disciplinary Venous Thromboembolism (VTE) Prevention Collaborative. *BMJ*. 2012;344:e3935. doi:10.1136/bmj.e3935
24. Pronovost PJ, Berenholtz SM, Needham DM. Translating evidence into practice: a model for large scale knowledge translation. *BMJ*. 2008;337:a1714. doi:10.1136/bmj.a1714
25. Haut ER, Lau BD, Kraenzlin FS, et al. Improved Prophylaxis and Decreased Rates of Preventable Harm With the Use of a Mandatory Computerized Clinical Decision Support Tool for Prophylaxis for Venous Thromboembolism in Trauma. *Arch Surg*. 2012;147(10):901-907. doi:10.1001/archsurg.2012.2024
26. Lau BD, Streiff MB, Hobson DB, et al. Beneficial “halo effects” of surgical resident performance feedback. *Journal of Surgical Research*. 2016;205(1):179-185. doi:10.1016/j.jss.2016.06.024

27. Elder S, Hobson DB, Rand CS, et al. Hidden Barriers to Delivery of Pharmacological Venous Thromboembolism Prophylaxis: The Role of Nursing Beliefs and Practices. *Journal of Patient Safety*. 2016;12(2):63. doi:10.1097/PTS.0000000000000086
28. Lau BD, Shaffer DL, Hobson DB, et al. Effectiveness of two distinct web-based education tools for bedside nurses on medication administration practice for venous thromboembolism prevention: A randomized clinical trial. *PLOS ONE*. 2017;12(8):e0181664. doi:10.1371/journal.pone.0181664
29. Owodunni OP, Lau BD, Streiff MB, et al. What the 2018 ASH venous thromboembolism guidelines omitted: nonadministration of pharmacologic prophylaxis in hospitalized patients. *Blood Advances*. 2019;3(4):596-598. doi:10.1182/bloodadvances.2018030510
30. Haut ER, Aboagye JK, Shaffer DL, et al. Effect of Real-time Patient-Centered Education Bundle on Administration of Venous Thromboembolism Prevention in Hospitalized Patients. *JAMA Network Open*. 2018;1(7):e184741. doi:10.1001/jamanetworkopen.2018.4741
31. Popoola VO, Lau BD, Shihab HM, et al. Patient Preferences for Receiving Education on Venous Thromboembolism Prevention – A Survey of Stakeholder Organizations. *PLOS ONE*. 2016;11(3):e0152084. doi:10.1371/journal.pone.0152084
32. Lau BD, Haider AH, Streiff MB, et al. Eliminating Healthcare Disparities Via Mandatory Clinical Decision Support: The Venous Thromboembolism (VTE) Example. *Med Care*. 2015;53(1):18-24. doi:10.1097/MLR.0000000000000251
33. Owodunni OP, Haut ER, Shaffer DL, et al. Using electronic health record system triggers to target delivery of a patient-centered intervention to improve venous thromboembolism prevention for hospitalized patients: Is there a differential effect by race? *PLOS ONE*. 2020;15(1):e0227339. doi:10.1371/journal.pone.0227339
34. Owodunni OP, Lau BD, Wang J, et al. Effectiveness of a Patient Education Bundle on Venous Thromboembolism Prophylaxis Administration by Sex. *Journal of Surgical Research*. 2022;280:151-162. doi:10.1016/j.jss.2022.07.015
35. Haut ER, Owodunni OP, Wang J, et al. Alert-Triggered Patient Education Versus Nurse Feedback for Nonadministered Venous Thromboembolism Prophylaxis Doses: A Cluster-Randomized Controlled Trial. *Journal of the American Heart Association*. 2022;11(18):e027119. doi:10.1161/JAHA.122.027119
36. Haut ER, Owodunni OP, Shaffer DL, McQuigg D, Samuel D, Hobson DB, Kraus PS, Wang J, Webster KLW, Kantsiper M, Harris JE Jr, Holzmueller CG, Varasteh Kia M, Streiff MB, Lau BD. Implementing a Patient-Centered Education Bundle to Improve Venous Thromboembolism Prevention. *JAMA Surg*. 2025 Oct 8:e254136. doi: 10.1001/jamasurg.2025.4136. Epub ahead of print. PMID: 41060638; PMCID: PMC12509082.
37. Owodunni OP, Lau BD, Shaffer DL, et al. Disseminating a patient-centered education bundle to reduce missed doses of pharmacologic venous thromboembolism (VTE) prophylaxis to a community hospital. *Journal of Patient Safety and Risk Management*. 2021;26(1):22-28. doi:10.1177/2516043520969324
38. Malerba SA, Fumagalli RM, Ay C, et al. Availability of medical and endovascular therapies for venous thromboembolism: a global survey for World Thrombosis Day. *Journal of Thrombosis and Haemostasis*. 2024;22(1):255-262. doi:10.1016/j.jth.2023.10.002
39. Nguyen TTT, Tong HT, Nguyen HTL, Nguyen TD. A Call to Action for Anticoagulation Stewardship to Address Suboptimal Thromboprophylaxis Practices for at-Risk Non-Orthopedic Surgical Patients in Vietnam: An Explanatory Sequential Mixed-Methods Study. *Vasc Health Risk Manag*. 2025;21:305-326. doi:10.2147/VHRM.S505100
40. Chahal R, Alexander M, Yee K, et al. Impact of a risk-stratified thromboprophylaxis protocol on the incidence of postoperative venous thromboembolism and bleeding. *Anaesthesia*. 2020;75(8):1028-1038. doi:10.1111/anae.15077
41. Lau BD, Shaffer DL, Kraus PS, et al. Prevention of Hospital-Associated Venous Thromboembolism: A Road Map to Defect-Free Care. *Quality Management in Healthcare*. 2023;32(4):281. doi:10.1097/QMH.0000000000000436