




KOSOVA JOURNAL OF SURGERY

TRAUMA AND CRITICAL CARE SURGERY UPDATE: EXPANDING THE EVIDENCE — PART II



Milestones and Key Figures in the Development of Damage Control Surgery

**Volume 10, Issue 1, Part II:
Trauma and Critical
Care Surgery Update:
Expanding the Evidence
April 2026
ISSN: 3027-5008 (Online)
ISSN: 3027-5016 (Print)**

VEDAT ELJEZI: State of the Art Review: Extracorporeal Membrane Oxygenation (ECMO): A Life-Saving Therapy And Its Challenges of Implementation In Emerging Healthcare Systems

NOAH ANTES, HUNTER ROW, THOMAS HALDIS, CORNELIUS DYKE: Transcatheter Aortic Valve Replacement for Severe Aortic Stenosis: A Review and Special Considerations

LULEJETA ATLANTIDA LATIFI, RIFAT LATIFI: Damage Control Surgery: An Update

WADE HOPPER, MENTOR AHMETI: State of the Art Interventions in Geriatric Trauma

DANIEL SCHEESE, TRAVIS SULLIVAN, JEFFREY HAYNES, EDGAR RODAS: Navigating the Challenges of Managing Pediatric Trauma: Current Trends, Evidence, and Practical Pathways

BELINDA DE SIMONE, FAUSTO CATENA, RIFAT LATIFI: Venous Thromboembolism Prophylaxis in Trauma and Critically Ill Patients: A State-of-the-Art Evidence-Based Review

OLUWAFEMI P. OWODUNNI, ELLIOTT HAUT: Implementing Sustainable Systems for Achieving Excellence in Venous Thromboembolism Prevention

Blunt Polytrauma in Two Pediatric Patients: Surgical vs. Conservative Management – A Two-Case Study

Shaban Memeti¹, Haris Sulejmani², Marjan Kamiloski¹, Rexhep Selmani³,
Qemal Rushiti³, Natalija Cokleska¹, Gani Ceku⁴, Defrim Kocinaj⁴

¹University Clinic of Pediatric Surgery, Skopje, North Macedonia; Faculty of Medicine, Ss. Cyril and Methodius University in Skopje, North Macedonia

²Faculty of Medicine, Ss. Cyril and Methodius University in Skopje, North Macedonia

³Faculty of Medicine, Ss. Cyril and Methodius University in Skopje, North Macedonia; University Clinic for Digestive Surgery, Faculty of Medicine, Ss. Cyril and Methodius University in Skopje, North Macedonia

⁴Clinic of Pediatric Surgery, University Clinical Centre of Kosovo, Prishtina, Kosovo

Corresponding Author:

HARIS SULEJMANI

Faculty of Medicine, Ss. Cyril and Methodius University in Skopje, North Macedonia

Email: sulejmani.haris@hotmail.com

Abstract

Background: Pediatric blunt abdominal trauma presents complex diagnostic and therapeutic challenges. This case series highlights two pediatric patients with polytrauma from road traffic accidents, each with distinct intra-abdominal injuries and differing management strategies.

Case Presentation: Case 1 involved a 7-year-old female with pulmonary contusion, facial lacerations, splenic rupture, and bowel perforation. Case 2 involved a 13-year-old female presenting with macrohematuria and left flank pain after ejection from a vehicle; imaging revealed a Grade III/IV renal laceration with retroperitoneal hematoma and minor pleural effusion.

Intervention: Case 1 underwent emergency

laparotomy with splenectomy and segmental bowel resection, followed by intensive monitoring and antimicrobial therapy. Case 2 was managed conservatively with intravenous fluids, antibiotics, and blood transfusion.

Outcome: Both patients recovered without major complications. Case 1 had a shorter hospital stay following definitive surgical control. Case 2 required prolonged observation due to the risk of delayed renal complications but achieved complete recovery.

Conclusions: Hemodynamically unstable patients with splenic rupture and bowel perforation require urgent surgery, while stable patients with high-grade renal injuries can be managed non-operatively, emphasizing individualized, stability-based care.

Keywords: pediatric trauma, blunt abdominal injury, polytrauma, splenic rupture, non-operative management, emergency surgery

Introduction

Blunt abdominal trauma (BAT) remains one of the most frequent and challenging scenarios in pediatric trauma care. Due to anatomical differences such as proportionally larger solid organs and thinner abdominal walls children are more prone to solid organ injuries (SOIs) than adults¹. The spleen, liver, and kidneys are the most commonly affected organs, with management strategies varying according to injury severity and institutional protocols².

Operative interventions such as splenectomy were once common, but advances in imaging and clinical practice have shifted care toward non-operative management (NOM) in hemodynamically stable children, as endorsed by APSA³⁻⁴. Renal trauma care has evolved similarly, with growing reliance on conservative treatment and selective angioembolization in stable patients⁶.

Previous reports from our center have described successful management of other pediatric abdominal emergencies, including urgent echinococcosis surgery⁷ and splenic torsion due to wandering spleen⁸, underscoring the importance of timely diagnosis and individualized treatment. Similar observations have also been reported in regional pediatric surgical literature⁹.

This case series compares two pediatric patients with similar mechanisms of injury but different intra-abdominal

injuries and management approaches, highlighting individualized decision-making.

Case Presentations

These two cases illustrate two different management approaches in pediatric abdominal trauma, aligning with current evidence and guidelines.


Case 1

A previously healthy 7-year-old female was brought to the emergency department after being struck by a car while crossing the street, sustaining facial lacerations, extremity excoriations, and abdominal trauma. She was hypotensive (85/50 mmHg) and tachycardic (125 bpm), consistent with hypovolemic shock.


Chest CT demonstrated pulmonary contusion with a minimal left hemothorax. Abdominal CT revealed splenic rupture with perisplenic hematoma (Figure 1). Laboratory testing showed significant anemia (87 g/L), leukocytosis ($17.2 \times 10^9/L$), elevated CRP (260 mg/L), and hypoalbuminemia (28 g/L).

Emergency laparotomy confirmed splenic rupture and descending colon perforation, and she underwent splenectomy and segmental bowel resection with primary anastomosis. Following surgery and source control, CRP levels (Graph 1) declined rapidly, hemoglobin improved after resuscitation, and all laboratory parameters progressively normalized. She was discharged in good condition after 5 days.

Figure 1: Contrast-enhanced CT findings in a pediatric patient with blunt abdominal trauma.
(A) Coronal CT demonstrating a large left renal laceration with surrounding perirenal hematoma (arrows).
(B) Axial CT image showing splenic rupture with heterogeneous splenic parenchyma (arrow).



Graph 1: Serial trends of C-reactive protein and hemoglobin in two pediatric trauma patients managed surgically and conservatively.



Case 2

A 13-year-old female passenger was ejected through a lateral window during a high-speed collision. She presented alert and hemodynamically stable (BP 100/50 mmHg, HR 87 bpm) with left flank pain, facial excoriations, and macrohematuria.

Abdominal CT demonstrated a Grade III/IV left renal laceration with perinephric hematoma and retroperitoneal fluid. Thoracic CT revealed mild pulmonary contusion and a small left pleural effusion (200 mL). Laboratory results showed anemia (105 → 91 g/L), elevated CRP (238 mg/L), and hypoalbuminemia (28 g/L).

She was managed non-operatively with intravenous fluids, analgesia, meropenem, ciprofloxacin, transfusion of one unit of packed red blood cells and plasma, tetanus prophylaxis, and gastroprotection. CRP (Graph 1) declined gradually, reflecting slower resolution of renal parenchymal inflammation, and hemoglobin stabilized after transfusion. She was discharged in good condition after 18 days.

Although both children sustained blunt abdominal trauma in motor vehicle accidents, their courses diverged due to differences in injury severity and physiological response. Patient 1 was unstable with splenic rupture and bowel perforation, requiring urgent laparotomy, resulting in rapid postoperative clinical and laboratory improvement. Patient 2, despite having a high-grade renal laceration, remained stable and successfully underwent conservative management with prolonged observation and

serial laboratory monitoring. Both recovered well, illustrating the effectiveness of stability-based, individualized treatment strategies in pediatric polytrauma (Table 1).

Discussion

The trend toward non-operative management (NOM) in pediatric abdominal trauma has significantly reduced the need for surgery while maintaining excellent outcomes in appropriately selected patients. Data from Bagwell et al. show that up to 78% of children with splenic trauma were successfully treated without surgery, with splenectomy rates markedly lower at pediatric trauma centers compared to adult centers (7.8% vs. 15.7%).¹ This underscores the importance of specialized pediatric care and the influence of institutional experience on surgical decision-making.

Similarly, liver injury grading based solely on imaging does not reliably predict clinical outcomes, reinforcing that hemodynamic stability should guide management rather than radiologic severity alone.¹ Current APSA-endorsed guidelines recommend tailored approaches based on vital signs, transfusion needs, and signs of ongoing bleeding.⁴

Renal trauma, while less frequent, follows a similar trajectory. Fernández-Ibieta emphasizes the value of four-phase contrast-enhanced CT in diagnosis and highlights the success of NOM, even in high-grade injuries, when patients remain stable.⁶ The growing role of interventional radiology, particularly selective angiobolization, has further reduced the need for operative

Table 1: Comparative summary of clinical characteristics and management in two pediatric patients with blunt abdominal trauma

Category	Patient 1	Patient 2
Age	7 years	13 years
Mechanism of Injury	Road traffic accident (RTA)	Blunt trauma (RTA, ejection)
Renal Injury	None	Grade III/IV renal laceration
Major Injuries	Splenic rupture, bowel perforation, pulmonary contusion	Renal laceration, pulmonary contusion, pleural effusion
Hemodynamic Stability	Unstable	Stable
Surgical Intervention	Yes	No
Management Approach	Emergency laparotomy (splenectomy, bowel resection)	Conservative (fluids, antibiotics, transfusion)
Hospital Course	ICU → Surgery → Step-down	Ward observation → Discharge
Length of Stay	5 days	18 days
Discharge Outcome	Good condition, transferred	Good condition, discharged

management in pediatric solid organ injuries by offering targeted hemostasis while preserving organ function.

Organ injury grading, such as the American Association for the Surgery of Trauma (AAST) classification, remains an important reference for assessing injury severity but should always be interpreted in clinical context. In this study, Case 1 represented a Grade V splenic injury with associated hollow viscus perforation, while Case 2 sustained a Grade III–IV renal injury. These grades correlate with the observed management strategies, where hemodynamic stability rather than grade alone determined treatment choice.

Access disparities remain a significant challenge. As Nance et al. found, more than 17 million U.S. children lack access to a verified pediatric trauma center within a 60-minute transport window, particularly in rural areas.⁵ This gap contributes to variability in care, including higher operative rates in centers unfamiliar with pediatric-specific trauma protocols.

Current reviews stress the importance of minimizing unnecessary interventions. Gates et al., through a systematic review, concluded that routine follow-up imaging, prophylactic embolization, and prolonged hospital stays offer little benefit in stable patients,⁴ while Lyttle et al. advocate reduced activity restrictions and shorter hospital stays in line with updated protocols prioritizing physiologic recovery over injury grade.⁵

While NOM is preferred for most solid organ injuries, hollow viscus injury remains a major indication for urgent laparotomy in pediatric blunt abdominal trauma.¹

Early recognition is essential, as delays in surgical intervention increase morbidity.

In summary, the shift toward non-operative management (NOM) has improved outcomes and optimized resource utilization in pediatric trauma care. Wider implementation of these strategies and equitable access to specialized pediatric centers are essential for consistent, evidence-based treatment. Surgery remains indispensable in hemodynamically unstable patients, in cases of ongoing hemorrhage, or when hollow viscus injury is suspected.

In our opinion, if Case 1 had sustained an isolated splenic rupture without associated bowel perforation or hemodynamic instability, non-operative management could have been considered, consistent with current APSA guidelines.

This study is limited by its descriptive nature and small sample size, which limits generalizability but provides valuable insight into individualized decision-making in pediatric trauma management.

Conclusion

This case series demonstrates that hemodynamically unstable patients with splenic rupture and bowel perforation require urgent surgical intervention, while stable patients with high-grade renal injuries can be safely managed non-operatively. A multidisciplinary approach, involving pediatric surgeons, radiologists, and critical care teams, remains essential for optimizing outcomes in pediatric blunt abdominal trauma.

Author Contributions

Shaban Memeti – Study conception and design, surgical management of patients, critical revision of the manuscript. Haris Sulejmani – Literature review, drafting of the manuscript, preparation of tables and figures.

Marjan Kamiloski – Surgical management of patients, acquisition of clinical data, manuscript editing.

Rexhep Selmani – Surgical consultation, contribution to case analysis, manuscript review.

Qemal Rushiti – Data collection, coordination between institutions, case documentation.

Natalija Cokleska – Data analysis, manuscript proofreading, and editing for clarity.

Gani Ceku – Case identification, literature review support, preparation of figure legends.

Defrim Kocinaj – Clinical data acquisition, assistance in drafting the discussion, manuscript review.

Abbreviations

BAT – Blunt abdominal trauma

SOI – Solid organ injury

NOM – Non-operative management

APSA – American Pediatric Surgical Association

CRP – C-reactive protein

ICU – Intensive care unit

RTA – Road traffic accident

CT – Computed tomography

Acknowledgments

None.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of Interest

The authors declare that they have no conflict of interest.

Ethical Approval

Ethical approval for this study was obtained from the Institutional Review Board of the Pediatric Surgery Clinic in Skopje, North Macedonia. All data were anonymized in accordance with institutional guidelines.

Consent for Publication

Written informed consent was obtained from the legal guardians of the patients whose imaging is included in this study. All identifying information has been removed to protect patient privacy.

References

1. Bagwell C. Management of blunt abdominal trauma in children. *AAP Grand Rounds*. 2002;8(1):8–9. Available from: <https://doi.org/10.1542/gr.8-1-8-a>

2. Notrica DM. Pediatric blunt abdominal trauma: current management. *Curr Opin Crit Care*. 2015;21(6):531–7. Available from: <https://doi.org/10.1097/MCC.0000000000000249>

3. Gates RL, Price M, Cameron DB, et al. Non-operative management of solid organ injuries in children: An APSA Outcomes and Evidence Based Practice Committee systematic review. *J Pediatr Surg*. 2019;54(8):1519–26. Available from: <https://doi.org/10.1016/j.jpedsurg.2019.01.012>

4. Nance ML, Carr BG, Branas CC. Access to pediatric trauma care in the United States. *Arch Pediatr Adolesc Med*. 2009;163(6):512–8. Available from: <https://doi.org/10.1001/archpediatrics.2009.65>

5. Lyttle BD, Williams RF, Stylianos S. Management of pediatric solid organ injuries. *Children (Basel)*. 2024;11(6):667. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC11202015/>

6. Fernández-Ibieta M. Renal trauma in pediatrics: a current review. *Urology*. 2018;118:45–50. Available from: <https://doi.org/10.1016/j.urology.2017.09.030>

7. Memeti S, Sulejmani H, Risteski T, Cokleska-Shuntov N, Racaj A, Ljumani-Bakui N, et al. Pediatric Echinococcosis: diagnosis, urgent surgical intervention, and treatment – case report. *Arch Public Health*. 2024. Available from: <https://journals.mk/index.php/aph/article/view/6150>

8. Memeti S, Kuci S, Ristevski T, Todorovic L, Kamilovski M, Krsteska B, et al. Clinical Management and Surgical Outcomes of Wandering Spleen with Splenic Torsion in Pediatric Patients: A Case Report. *Albanian J Trauma Emerg Surg*. 2025;9(1). Available from: <https://journal.astes.org.al/AJTES/index.php/AJTES/article/view/441>

9. Hyseni G, Kotorri B, Statovci S, Ademaj I, Hyseni F, Hyseni N. Large splenic cyst in a child: Case presentation. *Kosova J Surg*. 2023;7(1):68–71. Available from: https://koscs.org/wp-content/uploads/2023/03/Nexhmi-Hyseni_merged.pdf